# **Understanding Eastern Himalayan Cryosphere Using Remote Sensing**

by

## Saurabh Kaushik 10PP18J15003

A thesis submitted to the
Academy of Scientific & Innovative Research
for the award of the degree of
DOCTOR OF PHILOSOPHY
in
(SCIENCE)

Under the supervision of

Dr. Tejpal Singh and Prof. P.K. Joshi



## CSIR-Central Scientific Instruments Organisation, Chandigarh



Academy of Scientific and Innovative Research AcSIR Headquarters, CSIR-HRDC campus Sector 19, Kamla Nehru Nagar, Ghaziabad, U.P. – 201002, India

> January 2023

#### CERTIFICATE

This is to certify that the work incorporated in this Ph.D. thesis entitled, 'Understanding Eastern Himalayan Cryosphere Using Remote Sensing", submitted by Saurabh Kaushik to the Academy of Scientific and Innovative Research (AcSIR) in fulfillment of the requirements for the award of the Degree of DOCTOR OF PHILOSOPHY in SCIENCE, embodies original research work carried-out by the student. We, further certify that this work has not been submitted to any other University or Institution in part or full for the award of any degree or diploma. Research material(s) obtained from other source(s) and used in this research work has/have been duly acknowledged in the thesis. Image(s), illustration(s), Figure(s), table(s) etc., used in the thesis from other source(s), have also been duly cited and acknowledged.

Name: Saurabh Kaushik

Date: 13/02/2023

Name: Prof. P.K. Joshi Date: 13/02/2023

(Signature of Student) (Signature of Co-Supervisor) (Signature of Supervisor)

Name: Dr. Tejpal Singh

Date: 13/02/2023

#### **STATEMENTS OF ACADEMIC INTEGRITY**

I, Saurabh Kaushik, a Ph.D. student of the Academy of Scientific and Innovative Research (AcSIR) with Registration No. 10PP18J15003 hereby undertake that the thesis entitled "Understanding Eastern Himalayan Cryosphere Using Remote Sensing" has been prepared by me and that the document reports original work carried out by me and is free of any plagiarism in compliance with the UGC Regulations on "Promotion of Academic Integrity and Prevention of Plagiarism in Higher Educational Institutions (2018)" and the CSIR Guidelines for "Ethics in Research and in Governance (2020)".

( Koushik

Signature of the Student

Date: 13/02/2023 Place: Chandigarh

It is hereby certified that the work done by the student, under my/our supervision, is plagiarism-free in accordance with the UGC Regulations on "Promotion of Academic Integrity and Prevention of Plagiarism in Higher Educational Institutions (2018)" and the CSIR Guidelines for "Ethics in Research and in Governance (2020)".

Signature of the Co-supervisor

Name: Prof. P.K. Joshi

Date: 13/02/2023 Place: New Delhi Signature of the Supervisor

Name: Dr. Tejpal Singh

Jen L

Date: 13/02/2023 Place: Chandigarh

#### **ACKNOWLEDGEMENTS**

Writing a PhD is considerably more than just the process of writing a manuscript, the last five years have been a remarkable experience. I am thankful to so many people who have been a source of consistent support, inspiration and help over the last five years. Big thanks to my two supervisors, Dr. Tejpal Singh and Prof. P.K. Joshi for keeping me motivated throughout my PhD duration, and sending me all over the world for conferences, training and exchange programs. Prof. P.K. Joshi and Dr. Tejpal Singh have always pushed me to bring the best out of me and walked me through all sorts of technical and institutional challenges. I am highly thankful to my supervisors for their unconditional support and foster my scientific career beyond my expectations.

I am grateful to Director CSIR-CISO, AcSIR-CSIO coordinator (Prof. Vinod Karar and Prof. Sanjeev Soni) and Head of Department (Mr. Amitava Das) for consistent support, motivation, and all the required infrastructural facilities. I extend my gratitude to Mr. Suneet Sapra, assistant manager AcSIR-CSIO for his consistent support in smoothly inducting every AcSIR formality. I am also highly thankful to the entire Team of 'Polar and Cold region' (Prof. Claudia, Dr. Andreas Dietz, Dr. Celia Baumhoer, Dr. Mariel Dirscherl, Jonas Köhler; Marius Philipp, Dr. Sebastian Rößler at Germany Aerospace Center (DLR), where I spent 18 months as an exchange Ph.D. student. The incredible working experience at DLR has resulted in some of the best work carried out throughout my PhD duration. Danke schön!

This doctoral work would not have been possible without the support of various institutions which provided valuable funds and promoted me to excel without hardship. I would acknowledge the funding support from Innovation in Science Pursuit for Inspired Research (INSPIRE) GoI. I acknowledge the support received from Deutscher Akademischer Austauschdienst (DAAD) for granting the binationally supervised doctoral research grant that funded my travel and stay in Germany for a period of 18 months. The equipment support received from the Sustainable Natural Resource Use in Arctic and High Mountainous Areas (SUNRAISE) and Urban Resilience and Adaptation for India and Mongolia (URGENT) projects, co-funded by the Erasmus+ Programme of the European Union has helped in efficiently completing the laboratory work, thus are duly acknowledged. I would also like to thank the National Aeronautics and Space Administration (NASA), Japan Aerospace Exploration Agency (JAXA) and European Space Agency (ESA) for the free of cost data provided by them to carry out my research work.

Special thanks to all my colleagues and friends (Nitin Mohan Sharma, Vikas Kumar, Vikas Shaw, Dr. Lalit Maurya, Neha, Parul, Priyanka, Tarandeep Singh, Dr. Yogesh

Kumar, Raman Kaushal, Lavlesh, Vishavpreet Singh, and Vipasha Sharma) at CSIR-CSIO who have always been there for productive discussion over tea breaks, extended lunches, and dinners. I also owe thanks to my friends (Ram, Deepanshu, Pankaj, KK, Neeraj, Arvind, Ganjendra, Chinu, Divanshu and Vishal) for filling all these years with all sorts of adventure and fun. The unforgettable trips and time with all helped me to go through all the stressful times.

Lastly not the least I wish to thank my family: their support has always been a driving force for pursuance of excellence. My parents (Mr. Subhash Chand and Mrs. Raj Rani) unconditionally supported me throughout my career and held every decision of mine. My siblings (Dr. Jitendra Kaushik, Mr. Gaurav Sharma and Mrs. Madhuri) always took care of my financial and emotional support, things won't be the same and smooth without their support. Above all, all these won't be possible without the blessings of 'Lord Shiva'.

Saurabh Kaushik

## **Table of Contents**

CE	RTIFICATEii
ST	ATEMENTS OF ACADEMIC INTEGRITYiii
AC	KNOWLEDGEMENTSiv
Tal	ole of Contentvi
Lis	t of Figurex
Lis	t of Tablexv
Ab	breviationsxvi
Ch	apter 11
1.	Introduction
1	.1. Background1
1.2	Relevance of Himalayan Cryosphere4
	1.2.1. Physiographic relevance4
	1.2.2. Socio-economic relevance
1.3	State of Himalayan Cryosphere8
1.4	Research Motivation9
1.5	Research gaps and Objectives
1.6	Structure of Thesis
Ch	apter 213
2.	A review of Himalayan cryosphere with especial emphasis on Eastern Himalaya 13
2	.1. Glacier Mapping14
	2.1.1. Rationale of the glacier mapping
	2.1.3. Remote Sensing of glacier extent
	2.1.4. Mapping of debris cover glacier
	2.1.5. Existing approaches and emerging area of research
	2.1.6. Errors associated with glacier mapping
	2.1.7. Recommendations
2	.2. Glacier Velocity43
2	.3. State of glacial lake mapping47

Chapter 3		51
3. A meth	odological framework for automated Supraglacial-debris extraction	51
3.1. De	ep learning for glacier mapping	52
3.2. In	put Data	53
3.2.1.	Sentinel 1data	55
3.2.2.	Sentinel 2 data	55
3.2.3.	Landsat 8 data	55
3.2.4.	ALOS DEM	56
3.2.5.	Training labels	56
3.2.6.	Training and testing sites	57
3.3. Mo	ethodology	58
3.3.1.	Pre-Processing and Data Preparation	58
3.3.2.	Deep Neural Network Architecture (SGDNet)	60
3.3.3.	Post Processing	62
3.3.4.	Accuracy Assessment	62
3.4. Re	sults	64
3.4.1.	Proposed Deep Artificial Neural Network (SGDNet)	64
3.4.2.	Test Site 1, 2, and 3	66
3.4.3.	Generating glacier dataset for Sikkim Himalaya	70
3.5. Di	scussion	74
3.5.1.	Testing different band combinations	74
3.5.2.	Classification Results and Comparison with Existing Methods	77
3.5.3.	Limitations and Future Work	80
3.5.4.	Summary	81
Chapter 4		82
-	dological framework for automated glacial lake extraction	
	ep learning for glacial lake mapping	
	put Data	
4.2.1.	Training labels	
4.2.2.	Training and testing sites	
	ethodology	
4.3.1.	Pre-Processing and Data Preparation	
4.3.2.	Deep Convolutional Neural Network Architecture (GLNet)	

4.3.3.	Post Processing	92
4.3.4.	Accuracy Assessment	93
4.3.5.	GLOF hazard assessment	94
4.4. R	esults	96
4.4.1.	Proposed Deep Convolutional Artificial Neural Network (GLNet)	97
4.4.2.	Classification results for test sites (1, 2, 3 and 4)	97
4.4.2.	Generating glacier dataset for Sikkim Himalaya	102
4.5. D	iscussion	105
4.5.1.	Testing different band combinations	106
4.5.2.	Methodological advances and comparison with existing methods	108
4.5.3.	Methodological limitations and future requirement	113
4.5.4.	Summary	115
Chapter 5		117
5. Spatio	-temporal glacier surface velocity estimation using time series sate	ellite data 117
5.1. Glac	ier surface velocity estimation	117
5.2. Inpu	t Data	118
5.2.1. I	andsat Data	118
5.2.2. \$	SRTM DEM	119
5.2.3. 8	Study site	120
5.3. Metl	nodology	122
5.3.1. (	Glacier velocity and topographic control estimation	122
5.3.2. U	Incertainty estimation	125
5.4. Resu	ılts	126
5.4.1. 5	Spatiotemporal variation in glacier velocity	127
5.4.2. I	mpact of glacier size and debris cover on velocity	130
5.4.3. I	mpact of slope on glaciers velocity	132
5.5. Disc	ussion	135
5.5.1. l	Inderstanding spatiotemporal variability in glacier velocity	135
5.5.2. (	Comparison with other studies	138
5.5.4. S	Gummary	142
Chapter 6		143
6. Synth	esis and outlook	143

6.1.	Summary and conclusive findings	143
6.2.	Future outlook	146
Refere	nces	150
Appendix		
List of	Publications	186

# **List of Figures**

Figure 1.1 Cryosphere's component and their timescale; adapted from IPCC - AR4 WGI
Chapter 4: Observations: Changes in Snow, Ice, and Frozen Ground2
Figure 1.2 The Electromagnetic Spectrum: The remote sensing sensor are designed to sense
the specific range of EMR (usually varies from visible (0.4mm) to microwave (1m)). Credit:
miniphysics.com3
Figure 1.3 The extent of the Himalaya and Karakoram, major relevance is shown. The
boundaries are taken from Bolch et al., (2012). The MODIS and SRTM DEM is used as base
map6
Figure 2.1 Overview of glacier mapping studies in Himalaya16
Figure 2.2 Reflectance spectra of different snow and ice surfaces (Hall and Martinec 1985). 26
Figure 2.3 (a) Generalized cross-section of a typical valley glacier showing glacier boundary
as a boundary between supraglacial and periglacial debris, Figure courtesy of (Shukla et al.,
2010) (b) Shows the spectral similarity between supraglacial debris (SGD) and periglacial
debris (PGD). VRK, valley rock; MID, mixed ice and debris
Figure 2.4 Results of some semi-automated approach for debris cover delineation using a
combination of multiple dataset. (a) Debris cover glacier boundary delineated via
semiautomated algorithms using multispectral imagery and DEM derived parameter (Paul,
Huggel, and Andreas 2004), (b) Debris cover glacier boundary delineated using synergistic
approach (Shukla et al., 2010), (c) Glacier boundary extracted using hybrid CNN+ Radom
Forest (RF) approach (Nijhawan, Das, and Balasubramanian 2018), (d) Overlay of glacier
boundary delineated using combination of optical and thermal data over high resolution
satellite imagery (Karimi et al. 2012), (e) glacier boundary obtained via semi-automated
approach, utilizing input from optical, thermal and surface curvature (Bhardwaj et al. 2014).
This Figure illustrates, most of the studies demonstrated a semi-automated approach for
debris cover delineation on very small geographical area (i.e. 1or two glaciers) which reduces
the involved complexity of glacial terrain
Figure 2.5 (A) Map of some significant glacier velocity studies carried out in the Himalaya
and Karakoram. This map does not include study reported by Dehceq et al. (2019) which
provides a regional assessment of glacier velocity concentrated only on large (>5 km²) glaciers.
Yellow box shows location of study area, (ISM - Indian Summer Monsoon). Inset map shows
location of Himalaya and Karakoram on world map. The boundary of the Himalayan region
is taken from Bolch et al. (2012)
Figure 3.1 Spatial distribution of the test (blue) and training (red) sites across the Himalaya
and Karakoram regions. The star denotes region, which was used as both training and testing
data. The Himalayan and Karakoram boundaries were taken from Nie et al. (2017)
Figure 3.2 Deep learning neural network (DNN) architecture used for delineation of
supraglacial debris (SGDNet). The numbers indicate the total number of unit (i.e., neurons) in
each layer. The input layer consists of 9 bands (B: blue, G: green, R: red, NIR: near-infrared,
SWIR: shortwave infrared, T: thermal, C: coherence, S: slope, and E: elevation). The output of
the developed DNN is D: debris or ND: non-debris
THE REVELOPER DIVINED D. REDITO OF IND. HOH-REDITO

Figure 3.3 Workflow for automated supraglacial debris mapping using remote sensing and
deep learning63
Figure 3.4 Loss and accuracy curve of proposed DNN architecture (SGDNet) for SGD
classification65
Figure 3.5 Supraglacial debris mapping in Sikkim, Eastern Himalaya using the proposed
algorithm. False-positives and false-negatives were mapped manually using reference data.
Red, white, green, and yellow boxes provide a contextual overview of the Figure 3.14A, 3.14B,
3.14C, and 3.14D66
Figure 3.6 Supraglacial debris mapping at test site 1 (Northwest Karakoram) using the
proposed algorithm. False-positives and false-negatives were mapped manually using
reference data67
Figure 3.7 Supraglacial debris mapping at test site 2 (Karakoram) using the proposed
algorithm. False-positives and false-negatives are mapped manually using reference data 68
Figure 3.8 Supraglacial debris mapping at test site 3 (Karakoram) using the proposed
algorithm. False-positives and false-negatives are mapped manually using reference data. The
red box provides a contextual overview of Figure 3.1369
Figure 3.9 Compiled glacier dataset of Sikkim Himalaya70
Figure 3.10 Comparison of glacier boundaries mapped in the present study with the available
glacier inventories. The glacier boundary delineated in the present study also overlaid on
google earth image for visual comparison
Figure 3.11 Comparison of glacier boundaries mapped in the present study with the available
glacier inventories. The glacier boundary delineated in the present study also overlaid on
google earth image for visual comparison
Figure 3.12 The loss and accuracy curve of proposed DNN for SGD classification using a
combination of remote-sensing data. A - Optical-SAR-TIR-S; blue, green, red, NIR, SWIR,
coherence, and slope, B - Optical-TIR-S-E; blue, green, red, NIR, SWIR, and slope, C - Optical-
SAR-TIR; blue, green, red, NIR, SWIR, SAR, and TIR75
Figure 3.13 Qualitative analysis of results obtained using different combinations of bands. A
- Optical-SAR-TIR-S-E (blue, green, red, NIR, SWIR, coherence, slope, and elevation), B - Opti-
cal-SAR-TIR-S (blue, green, red, NIR, SWIR, coherence, and slope), C - Optical-TIR-S-E (blue,
green, red, NIR, SWIR, and slope), D - Optical-SAR-TIR (blue, green, red, NIR, SWIR, SAR,
and TIR). The classification results obtained from the Optical-SAR-TIR-S-E layer showed
better dis-crimination of SGD from PGS. The contextual information of Figure is presented in
Figure 3.8
Figure 3.14 Visual comparison of the results derived by the proposed algorithm with the
available reference data. RGI-corrected data show a corrected glacier boundary on ~2015
satellite data by Herreid and Pellicciotti (2020). The RGI updated shows the minor corrections
to Herreid and Pellicciotti (2020) glacier boundary as per scene characteristics. Supraglacial
debris data exhibit the supraglacial debris extent by Herreid and Pellicciotti (2020), and DNN-
derived shows the results of the proposed algorithm. The contextual information of Figure
3.14 A, B, C, and D is presented in Figure 3.5
Figure 4.1 Location map of chosen training (red) and test (yellow) sites across the Himalaya.
The inset map exhibits the location of the Himalayan and Karakoram. The glacier lake

boundaries are taken from Chen et al. (2021). The Karakoram and Himalayan boundaries were
adopted from Nie et al. (2017)86
Figure 4.2 Overview of the proposed GLNet. The inputs to the architecture are 10 pre-
processed bands (i.e., B, G, R, NIR, SWIR, TIR, SAR coherence, slope, elevation, and NDWI).
The features of the input image are an extracted series of convolution layers (encoder) and a
fully connected layer (decoder) providing the final output90
Figure 4.3 The overall workflow of the proposed scheme92
Figure 4.4 Pairwise comparison matrix of factors affecting Glacial Lake Outburst Flood
(GLOF)95
Figure 4.5 Pairwise comparison matrix of factors affecting Glacial Lake Outburst Flood
(GLOF)95
Figure 4.6 Glacial Lake mapping over test site 1, Eastern Himalaya using the proposed GLNet.
The GLNet derived lake boundaries are compared with the reference data to estimate error of
commission (FP) and omission (FN). The red and yellow rectangles show a location of lakes
displayed in the Figure 4.14 and Figure 4.1598
Figure 4.7 Glacial Lake mapping over test site 2, Eastern Himalaya using the proposed GLNet.
The GLNet derived lake boundaries are compared with the reference data to estimate error of
commission (FP) and omission (FN). The red rectangles show a location of lakes displayed in
the Figure 4.1499
Figure 4.8 Glacial Lake mapping over test site 3, Eastern Himalaya using the proposed GLNet.
The GLNet derived lake boundaries are compared with the reference data to estimate error of
commission (false-positive) and omission (false-negative). The red rectangles show a location
of lakes displayed in the Figure 4.14
Figure 4.9 Glacial Lake mapping over test site 4, Eastern Himalaya using the proposed GLNet.
The GLNet derived lake boundaries are compared with the reference data to estimate error of
commission (false-positive) and omission (false-negative). The red rectangles show a location
of lakes displayed in the Figure 4.14
Figure 4.10 Glacial lake dataset generated over entire Sikkim Himalaya. The stream network
is extracted via STRM 30m DEM using hydrological tools in ArcGIS103
Figure 4.11 Comparison of generated glacier lake dataset of Sikkim Himalaya with existing
datasets (e.g., Chen et al., 2020 and Zhang et al., 2021). The comparison reveals supremacy of
the dataset generated in the present study104
Figure 4.12 Glacial Lake Outburst Flood hazard assessment in Sikkim Himalaya105
Figure 4.13 The results of the proposed GLNet on all three-test site; (a) correlation analysis of
GLNet derived lake boundaries and reference lake boundaries; (b) box plot showing size
distribution of mapped glacial lakes via GLNet106
Figure 4.14 Example of precise glacial lake mapping via proposed GLNet across all three test
sites. GLNet successfully mapped glacial lakes of varied appearance and size across all three
test sites. T1; refers to test site 1, respectively T2; test site 2, T3; test site 3 and T4; test site 4.
The location of these lakes are displayed in the Figure 4.6, 4.7, 4.8 and 4.9
Figure 4.15 Visual comparison of most commonly used method for glacial lake mapping. A)
represent the test site; B) represent result derived via NDWI thresholding method; C) exhibits
automated glacial lake extraction using DNN (SGDNet(Kaushik et al., 2022b)); and D) shows

the results of the proposed method (GLNet) for glacial lake mapping. Contextual overview of
Figure is provided in Figure 4110
Figure 4.16 Visual comparison between DeepLabV3+ and GLNet for glacial lake extraction.
The GLNet outperformed DeepLabV3+, as DeepLabV3+ proved to be insignificant for
classifying glacial lakes with varied lake turbidity (D and E)112
Figure 4.17 Major limitations of GLNet; a, b, d represent an example of wet ice which is falsely
predicted as lake over test site 1 and 2; (c) represent false prediction of shadow as glacial lake
and (e) and (f) shows false negative prediction of frozen lakes over test site 2
Figure 5.1 (B) Map of study area showing studied glaciers and topography. The glacier
boundaries are taken from Randolph Glacier Inventory Version 6 (RGI V6) and Advanced
Land Observing Satellite (ALOS) - Phased Array type L-band Synthetic Aperture Radar
(PALSAR) (12.5 m) DEM is used for displaying topography. Fastest flowing glacier (ID 2,
Zemu) and all other marked glaciers such as ID0 = South Lohnak, ID21 = Middle Lohnak, and
ID22= North Lohnak, provide contextual information for Figure 5.6 and 4.7. The red and black
box provides a contextual overview of Figure 5.4 and 5.5
Figure 5.2 Flow chart of overall methodology125
Figure 5.3 Trend in glacier velocity of Sikkim Himalaya. Linear trend, the deviance bar shows
the estimated uncertainty in velocity for the particular image pair (A), Box plot showing
heterogeneity in glacier velocity in the Sikkim Himalaya (B). The uncertainty associated with
particular image pair is given in table 3127
Figure 5.4 Spatiotemporal variability in glacier velocity across Sikkim Himalaya. A) Glacier
velocity estimated during 1994-1996 and B) Glacier velocity estimated during 2017-2018.
Notice clear slowdown in the Zemu glacier128
Figure 5.5 Temporal snapshots of glacier slowdown in the Zemu glacier. A) velocity
estimation during 1994-1996, b) velocity estimation during 2009-2010 and c) velocity
estimation during 2017-2018. The glacier velocity is derived from image pairs of Landsat using
COSI-Corr. Notice clear slowdown in up glacier
Figure 5.6 Impact of glacier size and debris cover on glacier velocity in the Sikkim Himalaya
during 1994-2020. A) Velocity variation among the different sizes of Glaciers. B) Velocity
variation among different debris cover extent glaciers. DCG; Debris Cover Glacier (debris
cover >25%), SDCG; Sparsely Debris Cover Glacier (debris cover >10 and <25%) and CIG;
Clean Ice Glacier (debris cover < 10%).
Figure 5.7 Impact of slope on glacier velocity in the Sikkim Himalaya. (A) Relationship
between slope and glacier velocity considering all glaciers. (B) Relationship between slope
and glacier velocity for medium and large glaciers.
Figure 5.8 Computed Glacier velocity and surface slope along the central flow line for the
selected glaciers (G0, G1, G2, G3, G18, G19, G21, G23 and G50) during (1994-2020). The glacier
velocity and surface are resampled at every 120 m
Figure 5.9 Comparison between glacier velocity estimated along central flowline and
including all pixels for selected glaciers in the study region. Glacier velocity estimated along
central flowline exhibits higher velocity signal
Figure S.1Illustration of algorithm workflow and post-processing steps. A – results obtained
via proposed Deep Neural Network (SGDNet), B - results obtained via first post-processing

step (if DNN result>0.5 and elevation>5500), $C$ – cleaning up of result using 5×5 median filter
and D - shows results after vectorization, the red polygons (<0.02) are removed from final
results
Figure S.2 Impact of shadow on classification accuracy of the proposed algorithm. A -
presence of shadow in a satellite image, B - classification of shadow as false positive in the
results obtained via proposed algorithm167
Figure S.3 Examples of generated label data. The generated label data comprised of glacial
lakes of different size, shape and appearance. The special emphasis is laid on the inclusion of
frozen and partly frozen lakes
Figure S.4 Examples of generated label data. The generated label data comprised of glacial
lakes of different size, shape and appearance. The special emphasis is laid on the inclusion of
frozen and partly frozen lakes171
Figure S.5 Loss and accuracy curve of the proposed GLNet
Figure S.6 Detailed analysis of proportion of wet ice and shadow in false positive area
including all four test sites
Figure S.7 Off glacier pixels consider for error estimation (2001-2002)177
Figure S.8 Off glacier pixels consider for error estimation (2017-2018)177
Figure S.9 Scatter plot of glacier size and average glacier velocity during 1994-2020. The larger
glaciers correspond to higher glacier velocity
Figure S.10 Glacier area distribution of the studied glaciers, 128 glaciers are having size <10
km² (Out of 133 glaciers)178
Figure S.11 Four fastest flowing glaciated regions (red rectangles). All these regions are
present in the clean ice zones with icefalls and crevasses. (Image Soure: Copernicus Sentinel
2; scene Id = L1C_T45RXL_A019221_20201110T044603179

## **List of Tables**

Table 1.1 Glacier Inventory for Himalaya-Karakoram (HK) region compiled	from various
studies as referred in the text (Azam et al., 2019)	9
Table 2. 1 Overview of some significant studies shown in Figure 1.17	
Table 2. 2 Summary of some significant glacier's studies carried out in the Indi	an Himalaya
using manual digitization method.	21
Table 2. 3 Summary of some significant studies reported glacier delineation usir	ıg supervised
classification	28
Table 2. 4 Overview of some significant glacier velocity studies shown in Figure	2.545
Table 3.1 List of Sentinel 2 Spectral bands with central wavelength and resolu	ation (Source
Sentinel-2 user guides).	56
Table 3.2 Results of the proposed algorithm on the test sites	69
Table 3.3 The comparison of the generated glacier dataset with available glacie	er inventories
	71
Table 3.4. Results of the proposed deep artificial neural network (SGDNet) m	odel using a
combi-nation of optical, thermal, coherence, and slope as input layers	74
Table 4.1 Results of the proposed GLNet on the chosen test sites.	101
Table 4.2 Comparison of glacial lake (area and number) reported in the preser	nt study with
recently published datasets.	102
Table 4.3 A sensitivity analysis for different weighting schemes	105
Table 5.1 List of Landsat satellite data used to estimate glacier velocity	119
Table 5.2 Estimated uncertainty in velocity for each Image pair and average	for the entire
period	126
Table 5.3 Statistical results of Sikkim glacier velocity trend analysis (Mann-Kenc	lall and Sen's
slope)	134
Table S.1 Data used in the present study	167
Table S.2 Results of the various used Deep Artificial Neural Network models on	the test data.
	169
Table S.3 Lake characteristics of the chosen training data	173
Table S.4 Data used in the present study	
Table S.5 SIV of individual glacier over the entire study period (1994-2020)	180

## **Abbreviations**

ASTER Advanced Spaceborne Thermal Emission and Reflection Radiometer

COSI-Corr Co-registration of Optically Sensed Images and Correlation

CNN Convolutional Neural Network

DEM Digital Elevation Model

DCNN Deep Convolutional Neural Network

DNN Deep Neural Network
ELA Equilibrium Line Altitude
ESA European Space Agency

ETM+ Enhanced Thematic Mapper Plus
GIS Geographic Information System
GLOF Glacial Lake Outburst Flood

GLNet Glacial Lake Net

GSI Geological Survey of India

ICIMOD International Centre for Integrated Mountain Development

InSAR Interferometric Synthetic Aperture Radar

NASA National Aeronautics and Space Administration

NCC Normalized Cross Correlation

MSS Multispectral Scanner

NDSI Normalized Difference Snow Index

NDVI Normalized Difference Vegetation Index

NIR Near Infrared

NDWI Normalized Difference Water Index

OLI Operational Land Imager

PGD Periglacial Debris
ReLU Rectified Linear Unit
SAR Synthetic Aperture Radar

SGD Supraglacial Debris
SGDNet Supraglacial Debris Net

SNR Signal to Noise Ratio

SOI Survey of India

SRTM Shuttle Radar Topography Mission

SWIR Shortwave Infrared
TM Thematic Mapper
TSL Transient Snowline







#### **Last Page**

The appended material is based on research carried out at the partner institution of URGENT Project, and has potentially utilised the equipment support, inputs based on course revised/developed and training programs (*lecture series, research seminar and webinars*) through the URGENT Project.

The document is part of thesis part of PhD/MSc/MA research work carried out at the Jawaharlal Nehru University. Purposefully limited pages are shared to avoid copyright and other issues. However, the full thesis can be shared on request.

The complete thesis can be obtained from Prof P K Joshi (<u>pkjoshi27@hotmail.com</u> or <u>pkjoshi@mail.jnu.ac.in</u>).